



ORAL MICROFLORA AND DENTAL HEALTH- A REVIEW

Monika J. Mahajani¹, Chetan Agarwal², Vishwas D. Acharya*³, Shweta G. Habbu⁴

¹Professor and Head, Department of Periodontics, Dr Hedgewar Rugna Sewa Mandal, Dental College and Hospital, Hingoli, Maharashtra, India.

²Private Practice, Pune, Maharashtra.

³Senior Lecturer, Department of Oral Medicine and Radiology, Dr Hedgewar Rugna Sewa Mandal, Dental College and Hospital, Hingoli, Maharashtra, India.

⁴Senior Lecturer, Department of Public Health Dentistry, Dr Hedgewar Rugna Sewa Mandal, Dental College and Hospital, Hingoli, Maharashtra, India.

***Correspondence for Author: Dr. Vishwas D. Acharya**

Senior Lecturer, Department of Oral Medicine and Radiology, Dr Hedgewar Rugna Sewa Mandal, Dental College and Hospital, Hingoli, Maharashtra, India.

Article Received on 18/09/2015

Article Revised on 10/10/2015

Article Accepted on 02/11/2015

ABSTRACT

With the respect to microbial flora, the oral cavity is one of the most densely populated sites of the human body. The environmental diversity of the oral cavity promotes the establishment of distinct microbial communities, such as supragingival plaque, subgingival plaque and tongue coating. The properties of the environment determine which microorganisms can occupy a site, while the metabolic activities of those microbial communities subsequently modify the properties of the environment. The mouth contains both distinct mucosal (lips, cheek, tongue, palate) and, uniquely, non-shedding surfaces (teeth) for microbial colonisation. Each surface harbours a diverse but characteristic microflora, the composition and metabolism of which is dictated by the biological properties of each site. The resident oral microflora develops in an orderly manner via waves of microbial succession (both autogenic and allogenic). The mouth contains both distinct mucosal (lips, cheek, tongue, palate) and, uniquely, non-shedding surfaces (teeth) for microbial colonisation. Each surface harbours a diverse but characteristic microflora, the composition and metabolism of which is dictated by the biological properties of each site. The resident oral microflora develops in an orderly manner via waves of microbial succession (both autogenic and allogenic).

KEY WORDS: Oral Micro-organisms, Dental caries, Dental plaque, Periodontal bacteria.

INTRODUCTION

The oral flora is one of the most ecologically diverse microbial populations known to man. From birth the oral cavity is exposed to many different micro-organisms present in the local & geographic environment.^[1-5] The diversity of microflora is due to the fact that the mouth is composed of a number of varied habitats supplied with a range of different nutrients. In dental plaque gradients develop, such as oxygen tension, pH, providing conditions suitable for the growth & coexistence of micro-organisms. In order to understand the etiology of many oral & dental diseases^[6] & to interpret the results of microbiological analyses of clinical specimens, a knowledge of the micro-organisms which comprise the resident flora of oral cavity is important.

Acquisition of the oral flora

Acquisition depends on the successive transmission of micro-organisms to the site of potential colonization. Initially, in the mouth, this is by passive contamination

from the mother, from food, milk & water, & from the saliva of individuals in the close proximity to the baby.

By 24 hrs after birth the first species have become established. Streptococci, particularly *S.salivarius*, which binds to epithelial cells are usually the first to colonize. In first three days :- *S.oralis*, *S.mitis*. The diversity of the streptococcal microflora increases with time, so that all infants are colonized by more than one species of streptococcus by one month of age.^[7-9]

Table 1: Percentage viable count^[3-9]

<i>Streptococcus</i>	1-3 days	2 weeks	1 month
<i>S.oralis</i>	41	24	20
<i>S.mitis</i>	30	28	30
<i>S.salivarius</i>	10	30	28
<i>S.sanguis</i>	4	3	2
<i>S.anginosus</i>	3	5	5
<i>S.gordonii</i>	1	2	4

The diversity of the pioneer oral community increases during the first few months of life, & several gram-negative anaerobic species appear. *P.melaninogenica* – most frequently isolated anaerobe (76%) in infants with a mean age of 3 months. The dominant flora of the oral cavity in the child before eruption of teeth is mainly facultative in nature. With the eruption of teeth there is an increase in anaerobic forms, such as *Leptotrichia*, *Spirochetes*, *Fusiform bacilli*, *Spiral forms* & *Vibrio*. Complete loss of the dentition causes a reversion of the microflora to a predominantly aerobic facultative type.^[10-13]

Table 2: Other isolated species were

Micro-organism	Occasionally found	Usually found
<i>Veillonella</i>	< 2 days	After 1 week
<i>Fusiform bacilli</i>	< 2 months	Before eruption of incisors, appear to increase during 4 th & 8 th month
<i>Peptostreptococcus anaerobicus</i>	---	>5 months

Table 3

	Micro-Organism demonstrated
After eruption of teeth	<i>S.sanguis</i>
Primary dentition, when molars erupt	<i>S.mutans</i>
Predentate infants with acrylate cleft palate obturators	<i>S.Mutans</i>

Anaerobic forms generally reappear with the wearing of dentures.

e.g--*S.sanguis* & *S.mutans* disappear in edentulous mouth but reappear with wearing of dentures.^[8-13]

Table 4:

	Bacterial type present
Before eruption of teeth	Facultative
After eruption of teeth	Increase in anaerobic forms
Edentulous mouth	Aerobic facultative (reversion of the microflora)
On wearing dentures	Anaerobic forms reappear
Neglected/diseased mouth	Anaerobic & proteolytic
Well kept mouth	Aerobic & facultative
Healthy individual age 70 yrs/over	Lactobacilli ; Staphylococci
Healthy individual age 80 yrs	Yeast

It has been demonstrated that microbial plaque, after removal from tooth surface, begins to regenerate with in minutes – 1 million organisms were reported to deposit on 1 cm² of clean tooth surface with in 5 minutes. There appears to be an upper limit to which the microflora may increase in the oral cavity.^[14,15]

Factors that control & limit the population of oral microflora^[1,4,8-12]

- 1) Flushing action of saliva :- 1-2.5g of bacterial cells are swallowed each day in saliva.
- 2) Action of tongue, the lips, & mucous membrane of cheeks – helps to remove micro-organisms from tooth surface.
- 3) Desquamated exfoliated epithelial cells are shed, & with them the micro-organisms are removed & swallowed along with saliva.
- 4) Tissue fluids originating from sub-mucosal capillaries pass in to healthy gingival crevice & help remove micro-organisms from this area.

Micro-organisms in the oral cavity probably represent 2 phases of the growth curve: accelerated & stationary phase. After deposition, organism that become firmly attached to tooth enamel, should multiply & form communities of microcolonies. When the micro-organisms have increased to a certain population density, competition for nutrients between micro-colonies of the same & different organisms may result in retardation of cell division & in the death of some cells & thus ultimately cause decline of growth.

Distribution of oral microflora^[4-11]

The population making up the resident microflora community of the oral cavity are not uniformly distributed throughout the mouth.

Lip & palate

The lips form the border between the skin microflora (predominantly staph, micrococci, corynebacterium) & the mouth. Facultative anaerobic streptococci comprise a large part of the microflora on the lips. *S.vestibularis* is recovered most commonly from the gutter between lower lip & gums. *Candida albicans* can colonize damaged lip mucosal surfaces in the corner of the mouth (angular cheilitis).

Palate: majority of bacteria are streptococci, actinomyces, neisseria, veillonella.

Candida are not regularly isolated from the normal palate although their prevalence does increase if dentures are worn, & the mucosa is infected with *C.albicans*.

Cheek

Streptococci are the predominant bacteria from the cheek, especially *S.mitis*. Spirochaetes & other motile organisms have been occasionally attached to buccal mucosa. Haemophilus are commonly isolated in relatively large numbers.

Tongue

The dorsum of tongue, with its highly papillated surface, has a large surface area & therefore supports a higher bacterial density than other mucosal surfaces (100 bacteria / tongue epithelial cell). Streptococci are the predominant bacteria – 40% of the cultivable microflora. (*S.salivarius* & *mitis* group predominating) *Stomatococcus mucilaginous* – found exclusively on tongue. Veillonella : 16% Gram-positive rods : 16% Neissera : 20%

Saliva

Although saliva contains up to 10⁸ micro-organisms, it is not considered to have a resident microflora. Swallowing ensures that bacteria cannot be maintained in the mouth by multiplication in the mouth. The organisms found are derived from GCF flow, chewing, & oral hygiene but the tongue is the major source. Levels of streptococci / lactobacilli – is an indicator of the caries susceptibility of an individual.

Teeth

The microbial community associated with teeth varies in composition at each tooth surface due to differences in the local environmental conditions. *S.mutans*, *S.mitis*, *S.anginosus* – found in highest number on teeth as have strong affinity for hard surfaces. Gingival crevice debris develops in an environment (free from saliva). Supragingival plaque growing at the gingival margin on teeth influences the development of subgingival plaque. Plaque at gingival margin will tend to hinder the outflow of gingival fluid, epithelial cells & blocks entrance of saliva into the crevice. This environment would tend to encourage the establishment of anaerobic bacteria & the formation of the calculus.

Factors influencing bacterial adherence in oral cavity^[12-15]

Host factors

- 1) Mucosal cell surfaces contain sialic acid which binds to streptococcus mitis.
- 2) Enamel pellicle consists of albumin, lysozyme, immunoglobulins, proline -rich peptides & mucins. These act as receptors for oral bacteria.
- 3) Acidic proline-rich proteins: These promote the adherence of *Actinomyces* spp, *S.mutans* & black pigmented anaerobes.
- 4) Minerals, such as Calcium & phosphate, may influence the rate of calculus formation & provide larger surface areas for plaque to accumulate.
- 5) Many bacterial adhesins are lectins which bind to carbohydrate receptors on a surface.

Saliva

- 1) Mechanical washing action helps prevent microbial overgrowth
- 2) Salivary enzymes may inhibit bacteria. Example: Lysozyme – destroys bacterial cell walls.
- 3) Buffering capacity stabilizes pH to about 6.7. This prevents overgrowth of micro-

organisms which require high/low pH for maximal growth.

- 4) Salivary immunoglobulins, especially secretory IgA, may obstruct microbial adherence.

Bacterial factors

- 1) Adhesins are proteins associated with surface fimbriae & fimbriae.
- 2) Lipoteichoic acid (in bacterial cell walls) may help attachment to glycoproteins in tooth pellicle.
- 3) Extracellular polymers, example: Glucosyl transferase enzymes produce glucans which play a primary role in adhesion & support adherence of other microbial species.
- 4) Some bacteria possess IgA proteases – allowing them to overcome the blocking action of secretory IgA at mucosal surfaces.
- 5) Bacterial enzymes may expose binding site. example : Neuraminidase

CONCLUSION

Clinicians need to be aware of the beneficial properties of the resident microflora, and their treatment strategies should be focussed on the control rather than the elimination of these organisms, especially in dental plaque. In the future, it may be feasible to target treatment more specifically at particular 'pathogens' (e.g. immunotherapy (53)), or more imaginative approaches could be used to prevent disease.

REFERENCES

1. Houte JV, et al., In vitro acidogenic potential and mutans streptococci of human smooth surface plaque associated with initial caries lesions and sound enamel, *J. Dent. Res.* 70 (12) (1991) 1497–1502.
2. Sansone C. et al., The association of mutans streptococci and non-mutans streptococci capable of acidogenesis at a low pH with dental caries on enamel and root surfaces, *J. Dent. Res.* 72 (2) (1993) 508–516.
3. Houte JV, Lopman J, Kent R. The final pH of bacteria comprising the predominant flora on sound and carious human root and enamel surfaces, *J. Dent. Res.* 75 (4) (1996) 1008–1014.
4. Moore LVH et al., Bacteriology of human gingivitis, *J. Dent. Res.* 66 (5) (1987) 989–995.
5. Tanner A. et al., Microbiota of health, gingivitis, and initial periodontitis, *J. Clin. Periodontol.* 25 (2) (1998) 85–98.
6. Marsh PD. Microbial ecology of dental plaque and its significance in health and disease, *Adv. Dent. Res.* 8 (2) (1994) 263–271.
7. Saito K. et al., Effects of glucose on formation of cytotoxic end-products and proteolytic activity of *Prevotella intermedia* and *Prevotella nigrescens*, *J. Periodontol. Res.* 36 (6) (2001) 355–360.
8. Washio J. et al., Hydrogen sulfide-producing bacteria in tongue biofilm and their relationship with oral malodor. *J. Med. Microbiol.* (in press).

9. Ratcliff PA, Johnson PW. The relationship between oral malodor, gingivitis, and periodontitis. A review, *J. Periodontol.* 70 (5) (1999) 485– 489.
10. Kolenbrander PE, London J. Adhere today, here tomorrow: oral bacterial adherence, *J. Bacteriol.* 175 (11) (1993) 3247– 3252.
11. Takahashi N, Yamada T. Acid-induced acidogenicity and acid tolerance of non-mutans streptococci, *Oral Microbiol. Immunol.* 14 (1) (1999) 43–48.
12. Li Y, Caufield PW. The fidelity of initial acquisition of mutans streptococci by infants from their mothers. *J Dent Res* 1995; 74: 681–5.
13. McCarthy C, Snyder ML, Parker RB. The indigenous oral flora of man. I. The newborn to the 1-year-old infant. *Arch Oral Biol* 1965; 10: 61–70.
14. Milnes AR, Bowden GH, Gates DRT. Normal microbiota on the teeth of preschool children. *Microb Ecol Hlth Dis* 1993; 6: 213–27.
15. Caufield PW, Cutter GR, Dasanayake AP. Initial acquisition of mutans streptococci by infants: evidence for a discrete window of infectivity. *J Dent Res* 1993; 72: 37–45.