



DETOXIFICATION OF PATULIN MYCOTOXIN USING BY SEAWEED EXTRACT OF BROWN ALGAE ASCOPHYLLUM NODOSUM

Dr. Shatha Ali Shafiq*, Rana H. Al-Shammari, Sabaa Taher, Shahrazad Najem Abdu-Allah and Lika Hamid Mahdi

Mustansiriyah University - College of Science / Dept. Biology.

*Corresponding Author: Dr. Shatha Ali Shafiq

Mustansiriyah University - College of Science / Dept. Biology.

Article Received on 15/09/2017

Article Revised on 06/10/2017

Article Accepted on 27/10/2017

ABSTRACT

The current study was focused on the ability of the foreign product seaweed extract of brown algae *Ascophyllum nodosum* to detoxifying of Patulin mycotoxin producing from *Penicillium expansum*. The results showed that seaweed extract of brown algae at concentrations (1.5 and 3) % v/v inhibited the vegetative growth of *Penicillium expansum* the inhibition percentage reached to 79% and 90 % at the concentrations (1.5,3) % v/v respectively in culture medium Glucose-Czapek's Apple broth (GCA). However the results of test the ability of two concentrations of seaweed extract of brown algae in Patulin detoxification from broth culture of (GCA) for 14 days of incubation showed completely detoxification of Patulin compared with control treatment 325.01 µg/ml.

KEYWORDS: Patulin – Detoxification - seaweed extract.

INTRODUCTION

Mycotoxins are natural secondary metabolites produced by filamentous fungi of kingdom fungi, more than 500 types of mycotoxins have been identified to present, which include the most commonly mycotoxins associated with food and feed that may be concern to consumer food safety: Aflatoxins, Ochratoxin A, Patulin and Trichothecenes.^[1] Among the most important mycotoxins was (Patulin) PAT is a β unsaturated lactone and soluble in water and most polar organic solvents – soluble, produced by a number of fungal species belonging to the genera *Penicillium*, *Aspergillus*, *Byssochlamys* and *Pacelomyces*.^[2] The main producer of PAT is the blue mold *Penicillium expansum*, which is considered as a wound pathogen.^[3] It is usually associated with fruits and vegetables especially in apple and its products.^[4,5] PAT was first isolated as an antimicrobial active principle during 1940, from *Penicillium griseofulvum*, during the 1960, PAT was reclassified as a mycotoxin which was toxic to both plants and animals.^[6]

Most countries have set up maximum admissible levels of PAT in processed products from apples such as apple juice, apple concentrate juice, which have been identified 50 µg/L as recommended by the Food and Drug Administration.^[7] PAT exhibits sever toxicoses to human and animals exposed to the toxin via consumption of contaminated products, it has been shown to have toxic effects and can harm the immune system and gastrointestinal tract. Also shows carcinogenic,

teratogenic, and mutagenic properties toward different mammalian cells including human cells.^[8,9,10,11]

Different physical and chemical methods have been established for elimination of fungi producing mycotoxins and contamination of food and feed with toxigenic fungi. Nevertheless, only few of these methods have been accepted for practical uses, a practical and effective method was needed to be developed for the detoxification of PAT contaminated agricultural commodities.^[12]

Biological method using antagonist microorganisms has received large attention in recent years as a promising alternative, which considers more safety in both human health and ecosystem. The term probiotic was currently referred to ingestive microorganisms associated with beneficial effects to humans and other animals and plants.^[13,14] Some strains of lactic acid bacteria group (LAB) and yeast *Saccharomyces cerevisiae* have been reported to be effective in removing PAT from contaminated fruits.^[15,16] The aim of present study included inhibition of growth of toxigenic fungus *P.expansum* and detoxification of Patulin mycotoxin using by Seaweed extract of brown algae *Ascophyllum nodosum* in vitro.

MATERIAL AND METHODS

Fungal isolates

Three fungal isolates of *P. expansum* isolated from dried apricot fruit collected from different markets in Iraq at

Baghdad city included Palastine street (P), Al-Sadr city (S) and (Z) in Al- Utaifiyya, were obtained from department of biology - college of Science / Mustansiriyah university.

Detection of PAT Toxigenic Fungi

Glucose-Czapek's Apple medium (GCA)^[17] was used to stimulate PAT toxigenic fungi, conical flasks containing 50 ml of sterilized GCA inoculated with two discs (5 mm in diameter) of 7 days old culture grown on PDA, flasks were incubated at temperatures 15°C for 14 days in order to select incubation periods for production of PAT. Fungal biomass was separated by filtration through Whatman No.1 filter paper, then the filtrate was re-filtered through a millipore filter (0.45 µm in diameter) to remove fungal spore, 10 ml of filtrate was washed with 20 ml of chloroform in separating funnel, shaken for ten min. the top layer of chloroform was filtrated then reduced at 45°C then remained residue was dissolved in 1ml acetonitrile and kept in a deep freeze until used for PAT detection by (HPLC) the amount of PAT was estimated in comparison with standard PAT through the following formula.^[18]

$$\text{PAT conc.} = \frac{\text{Peak area of sample}}{\text{Peak area of standard}} \times \text{Standard concentration}$$

The percentage of reduction of PAT was calculated using the formula:-

$$\% \text{ Reduction} = \frac{\text{conc. in control treatment} - \text{concentration in tested sample}}{\text{concentration in control treatment}} \times 100.$$

Seaweed Extract

Seaweed extract of brown algae *Ascophyllum nodosum* is obtained as a commercial foreign product from College of Agriculture -Baghdad university.

Suppression of PAT by Seaweed Extract

The method described by Shafiq and others^[19] to detoxifying of PAT Seaweed extract of brown algae *Ascophyllum nodosum* in two concentrations (1.5 and 3) % v/v were added to 50ml of GCA broth containing toxigenic fungi produce PAT (*P. expansum* isolated from Palastine street P) and incubated at 15°C for 14 days, control treatment involved GCA broth containing toxigenic fungi produce PAT only with no seaweed extract addition, all treatments performed in triplicate then PAT concentration in all treatments were estimated by HPLC and the percentage of Patulin reduction was calculated.

RESULTS AND DISCUSSION

Evaluation of Fungal toxigenic

Three isolates of *P. expansum* from Palastine street, Al-Sadr City and Al- Utaifiyya, were evaluated on GCA medium at 15°C for 14 days, for their ability to produce PAT. (Table.1)

Table 1: The quantitative estimation of PAT in dried apricot fruit samples collected from various local markets in Baghdad.

Fungal isolates	Con. of PAT. µg/ml
<i>P. expansum</i> /P	325.01
<i>P. expansum</i> /S	51.1
<i>P. expansum</i> /Z	79

In this respect the amount of PAT production by fungal tested isolates in this study agreed with McCallum and others^[20] who found substantial differences among *P. expansum* strains in terms of growth kinetics and PAT production in three different apple ciders. The results also compatible with^[17,19] who showed maximum production of PAT using same medium (GCA) and different isolates of *P. expansum* and they noted that GCA was found to be the best medium supporting maximum PAT production by *P. expansum* isolates.

Effect of Seaweed Extract on Growth of *P. expansum*

The anti-fungal activity of seaweed extract are represented by the percentage of inhibition between two concentrations (1.5,3) % v/v of seaweed extract were increased with the increasing of concentration, the percentage of inhibition was reached to 79% and 90% at the concentrations (1.5,3) % v/v respectively in culture medium GCA compared with control treatment.

This result agreed with the findings of many studies^[21,22,23] they screened the most active compounds in algae, biochemical analysis are being undertaken to determine the structure and nature of compounds responsible of the bioactivity of the extracts with high antibacterial activity. Not only the presence of a particular compound which makes these organisms, interesting but also their huge diversity and the possibility of not only harvesting them but also of growing them at different conditions, leading to an enrichment of some bioactive compounds.

Suppression of PAT by Seaweed extract

The results of the activity of two concentrations of Seaweed extract (1.5 and 3) % v/v in suppression of PAT produced by *P. expansum* in toxigenic fungal culture medium, revealed that both concentrations completely detoxification the toxin from the culture medium with Reduction percentage 100%, while it recorded 325.01 µg/ml in control treatment Table -2.

Table 2: The Efficiency of Seaweed extract in reduction of PAT in GCA broth.

Concentration of Seaweed extract % v/v	Conc. PAT (µg/ml)	Reduction %
0.0	325.01	0.0
1.5	0.0	100
3	0.0	100

The detoxification activity of Seaweed extract may be due to their bioactive compounds and their interaction

with the patulin biosynthetic pathway. Our finding agreed with Nezha and others^[24] how reported the ability of ethanolic extracts marine algae (*Cystoseira tamaricifolia*) at the concentration of 5% to inhibit Mycotoxins formation in *Aspergillus flavus* and also Shafiq^[25] was shown the ability of sea weed extract of brown algae *Ascophyllum nodosum* as a detoxifying agent against mycotoxin Zearalenone production.

REFERENCES

- Magan, N., D. Aldred and V. Sanchis, The role of spoilage fungi in seed deterioration in: Arora, D.K. (eds.), *Fungal Biotechnology in Agricultural, Food and Environ. Appl.*, 2004; 311-123.
- Drusch, S., S. Kopka and J. Kaeding, Stability of Patulin in a juice – like aqueous model system in the presence of ascorbic acid, *food chemistry*, 2007; 100(1): 192-197.
- Lai, L., M. Fuh and C. Shih, Detection of mycotoxin Patulin in apple juice. *J. Food and Drug Analysis*, 2000; 8: 85-96.
- Ritieni, A., Patulin in Italian commercial apple products. *J. Agri and food chemi*, 2003; 51(21): 6086-6090.
- Richard, J.L. Some major mycotoxins and their mycotoxicoses—an overview. *Int. J. Food Microbiol.*, 2007; 119(1–2): 3–10.
- Andersen, B., J. Smedsgaard and J.C. Frisvad, *Penicillium expansum*: consistent production of patulin, chaetoglobosins and other secondary metabolites in culture and their natural occurrence in fruit products. *J. Agricult. Food Chem.*, 2004; 52: 2421-2428.
- World Health Organization (WHO) and Food and Agriculture Organization (FAO) of the United Nation. Safely evaluation of certain mycotoxins in food: Aflatoxin M1, fumonisins, ochratoxin A, trichothecenes, deoxynivalenol, T-2 and HT-2. Prepared by the 56th meeting of the Joint FAO/WHO Expert Committee on food additives, WHO Food Additives Series 47 and FAO Food and Nutrition paper74, WHO Geneva, 2001; 701.
- Alves, I.; Oliveira, N.G.; Laires, A.; Rodrigues, A.S. and Rueff, J. Induction of micronuclei and chromosomal aberrations by the mycotoxin patulin in mammalian cells: role of ascorbic acid as a modulator of patulin clastogenicity. *Mutagenesis*, 2000; 15: 229–234.
- Bing Hui, L.; FenYih, Y. and Tinsuan, W. Evaluation of genotoxic risk and oxidative DNA damages in mammalian cells exposed to mycotoxin, Patulin and Citrinin. *Toxicol. Appl. Pharmacol.*, 2003; 1919: 255-263.
- Shafiq, SH. A. Using of Iraqi probiotic to detoxify Patulin in albino mice. *International Journal of Innovation and Applied Studies*, 2015; 11(2): 282-290.
- Schumacher, D.; Metzler, M. and Lehmann, L. Mutagenicity of the mycotoxin Patulin in cultured Chinese hamster V79 cells and its modulation by intracellular glutathione. *Archives of Toxicology*, 2005; 79: 110-121.
- Verma, J., S. Shukla and N. Mehta, Amelioration of cytotoxic effects of Aflatoxin by vitamin A: an in vitro study on erythrocytes. *Toxicol. In Vitro*, 2008; 15: 39-42.
- Food and Agricultural Organization of the United Nations and World Health Organization (FAO). "Probiotics in food: health and nutritional properties and guidelines for evaluation", 2002.
- Rijkers, G. T., W. M. de Vos, R. J. Brummer, L. Morelli, G. Corthier, P. Marteau, De Vos, B. Morelli, Corthier and Marteau, "Health benefits and health claims of probiotics: Bridging science and marketing". *British Journal of Nutrition*, 2011; 106(9): 1291-6.
- Dalie, D., M. Deschamps and F. Richard-Forget, Lactic acid bacteria-Potential for control of mould growth and mycotoxins: A review *Food control*, 2010; 21: 370-380.
- Wang, L., T. Yue, Y. Yam, Z. Wang, M. Ye and R. Cai, A new insight into the adsorption mechanism of Patulin by the heat in active lactic acid bacteria cells, *food control*, 2015; 50: 104-110.
- Hasan, H.A.H., Patulin and aflatoxin in brown rot lesion of apple fruits and their regulation. *World Journal of Microbiology and Biotechnology*, 2000; 16: 607-612.
- EEC, Official Journal of the European communities. L327/54, 1992: 13(11): 92.
- Shafiq, SH.A; Aboud, H.H; Ismael, S. J. Biological Degradation of Patulin Mycotoxin Producing from some Fungi in Dried Fruits, *Advances in Environmental Biology*, 2016; 10(11): 107-111.
- McCallum, J.L.; Tsao, R. and Zhou, T. Factors affecting Patulin production by *Penicillium expansum*. *J. Food Protect*, 2002; 65: 1937–1942.
- Yousif, D.Y. M.; Dwish, A.S. and Shafiq, Sh. A. Antifungal activity of algal *Spirogyra* sp. against fungal *Fusarium Oxysporum*. *World Journal of Pharmaceutical Research*, 2015; 4(1): 1620-1628.
- Peres, J.C.F.; Carvalho, L.R. and Gonçalves, E. Evaluation of antifungal activity of seaweed extracts, *Ciênc. agrotec.*, Lavras, 2012; 36(3): 294-299.
- Dermirel, z.; Yilmaz-Koz, F.; Yavasoglu, K.; Ozderir, G. and Sukatar, T. Antimicrobial and antioxidant activity of brown algae from the Aegean Sea. *J. Serb. Chem. Soc*, 2009; 74(6): 619–628.
- Nezha, S.; Mohamed, L.; Mohamed, F.; Khadija, F.Z. Inhibition of growth and mycotoxins formation in moulds by marine algae *Cystoseira tamariscifolia*, *African Journal of Biotechnology*, 2004; 3(1): 71-75.
- Shafiq, SH. A. Antagonistic activity of probiotic and sea weed extract against vegetative growth for some fungi and Zearalenone production, *World J. Pharmaceutical Research*, 2015; 4(1): 1577 – 1585.