



FUNCTIONAL OUTCOME IN DISTAL TIBIA FRACTURES BY MIPPO TECHNIQUE USING TIBIA LOCKING PLATE

Dr. Allen Tilak J. T.*¹ and Dr. M. R. Rajasekhar²

¹Junior Resident, Department of Orthopedics, Sree Balaji Medical College, Chromepet, Chennai.

²Professor, Department of Orthopedics, Sree Balaji Medical College, Chromepet, Chennai.

*Corresponding Author: Dr. Allen Tilak J. T.

Junior Resident, Department of Orthopedics, Sree Balaji Medical College, Chromepet, Chennai.

Article Received on 21/06/2018

Article Revised on 11/07/2018

Article Accepted on 02/08/2018

ABSTRACT

Introduction: Distal tibial fractures constitute 1-10% of all lower extremity fractures. Operative treatment is indicated for most of the fractures which allows early motion and avoids shortening and other complications associated with prolonged immobilization. Minimally invasive percutaneous plate osteosynthesis (MIPPO) is a technique which aims to reduce iatrogenic soft tissue injury and damage to bone vascularity, as well as preserves the osteogenic fracture hematoma. **Material and Methods:** In the present study, twenty four patients with extra articular tibia were operated with MIPPO with distal tibia medial locking plate over period of two years and the results were analyzed. **Results:** At the end of the our study of 24 patients treated 14(58.3%) patients had excellent outcome 5(20.8%) had good results, 4(16.7%) had fair outcome and 1(4.1%) had a poor result based on subjective result and 16(66.7) patients had excellent, 4(16.7%), and 2(8.3%) had poor result based on objective result according to scoring system. **Conclusion:** Extra articular distal tibia fracture can be effectively treated with MIPPO technique using medial distal tibial locking plate.

KEYWORDS: Distal tibia fractures; MIPPO, Distal plate tibia locking.

INTRODUCTION

Distal tibial fractures constitute 1-10% of all lower extremity fractures.^[1] Controversy and debate continues to surround the management of fractures of the distal tibia.^[2] While non-operative treatment is associated with joint stiffness in up to 40% of cases as well as shortening and rotational malunion in over 30% of cases.^[3] Operative treatment allows early motion, and avoids shortening and other complications associated with prolonged immobilization.^[4] Conventional ORIF requires extensive soft tissue dissection with consequent periosteal injury even in expert hands. Complications such as malunion, delayed union and infection are at higher rates with ORIF. Intramedullary nails often do not provide enough stability and external fixators which are usually used for preliminary reduction and stabilization is also associated with complications like pin tract infection, delayed union and malunion. Recently, the trend is towards use of a Locking compression plate for treatment of fractures of the distal part of the tibia.^[6,7] Compared with a conventional plate, a Locking compression plate imparts a higher degree of stability and provides better protection against primary and secondary losses of reduction and minimization of bone contact.^[8,9] Minimally invasive plate osteosynthesis (MIPO) is a technique which aims to reduce iatrogenic soft tissue injury and damage to bone vascularity, as well

as preserving the osteogenic fracture haematoma.^[10] This philosophy is especially applicable in the management of distal tibial fractures, owing to the vulnerable extraosseous-metaphyseal blood supply in the distal region of the tibia.^[11] The purpose of this study was to evaluate the functional outcome, duration of union, advantages and complication following use of MIPO with LCP for meta-diaphyseal fractures of distal tibia.

MATERIALS AND METHODS

We undertook a review of 24 patients treated by MIPO (Minimal Invasive Plate Osteosynthesis) for unstable fractures of the distal tibia in MVJ Medical College and Research Hospital between June 2013 to June 2015.

Inclusion criteria

1. Adults (aged more than 18 years) males and females.
2. Complex extraarticular meta-diaphyseal fractures of the lower third of tibia.

Exclusion criteria

1. Open fractures.
2. Intra articular fractures.
3. Immuno compromised patients.

The mean age of the study group was 40.8 years (Range 22-58). Out of 24 patients, 21 (87.5%) are male patients and 3 (12.5%) were female patients showing increased

male preponderance in view of travelling, working in fields and factories. In our study 17 patients (70.8%) sustained injury following RTA and 7 patients (29.1%) sustained injury following fall from height. The fracture pattern was classified based on Rudie and Allgower classification for fractures of distal tibia. Of the 24 cases studied, 5 are A1, 8 are A2, 11 are A3. 16 out of 24 cases studied had an associated fracture of the lower third of fibula and 2 had upper third fibula fracture. Local examination of the injured extremity revealed swelling, deformity and loss of function. Palpation revealed abnormal mobility and crepitus at the fracture site. Distal neurovascular status was assessed based on posterior tibial artery and dorsalispedis artery pulsations, capillary filling, local temperature, pallor and paraesthesia. Antero-posterior and lateral radiographs of the affected leg along with ankle were taken and the fracture patterns were classified. Immobilization of the limb was done with an above knee Plaster of Paris slab till definitive fixation with locking compression plate done. All the cases were taken for early primary fixation with locking compression plate. 3 cases were delayed for 10 days due to swelling and those cases were operated after appearance of wrinkle sign.

Surgical Technique: Medial approach is most commonly used for the Mipo technique. Plate inserted from distal to proximal through epiperiosteal tunnel between periosteum and intact soft tissue.

Fracture is reduced by indirect maneuvers using ligamentotaxis and directly by percutaneous reduction forceps. 3-5 cm of slightly curved skin incision on medial aspect of distal tibia from the tip of medial malleolus. Incision carried out straight through subcutaneous fat without raising flaps. Great saphenous vein and nerve are held anteriorly ORIF of fibula was

part of first stage of fixation, which aids reduction of tibia. Epiperiosteal tunneling made towards the diaphysis by blunt tip of plate or tunneling instrument. Plate is inserted from distal to proximal on anteromedial surface using drill sleeve as plate manipulator. Fracture is by passed and plate is first adjusted to periarticular area and locking screws inserted in one the distal holes just above ankle joint approximating the plate to bone which prevents overlying soft tissue irritation. Fracture reduced by closed reduction maneuvers. Using another drill sleeve as manipulator at proximal end, the plate is fixed using locking cortical locking screw. Rest of the screws are inserted under c arm guidance using stab incisions atleast three on either side. Tourniquet was removed, hemostasis was secured and incisions closed with 2'0 nylon.

Post-operative regimen: Active toe movements after recovery from anaesthesia was started. Intravenous antibiotic regimen was continued for 3 days after the surgery. Patient was put on non weight bearing mobilization. Suture or staple removal was done at 10th to 14th post operative day. Patient progressed to full-weight bearing after x-ray showed evidence of callus formation.

Follow up: Routine follow up visits were scheduled at intervals of 4 to 6 weeks to assess the radiological union.

RESULTS

All the fractures united with an average of 16 weeks. There were 2 delayed union with 20 weeks signs of radiological callus formation. Fractures of 10 (41.7%) patients united in 14. weeks, 6 (25%) patients united in 16 weeks, 6 (25%) patients united in 18 weeks and 2 (8.3%) patients united in 20 weeks. All cases fibula fracture healed without complications.

OBJECTIVE AND SUBJECTIVE CRITERIA

Table. 1: Objective Criteria.

Rating	Ankle/ subtalar motion	Tibiotalar alignment	Tibial shortening	Chronic swelling	Equines Deformity
Excellent	>75% normal	Normal	None	None	None
Good	50-75%	Normal	None	Minimal	None
fair	25-50%	<50%angulation	<1cm	Moderate	None
poor	<25%	>50%angulation	>1cm	Severe	present

Table. 2: Subjective Criteria.

Rating	Pain	Return to Work	Recreational Activity	Limited Walking	Pain Medication	Limp
excellent	None	Same	Normal	No	None	None
Good	Mild	Same	Mild modification	No	None	None
Fair	Moderate	Moderate	Significant modification	Yes	Non narcotic	Occasional
poor	severe	Unable	None	yes	Narcotic	Yes

Table. 3: Objective Results.

Results	No. of cases	Percentage
Excellent	16	66.7
Good	4	16.7
Fair	2	8.3
Poor	2	8.3

Table. 4: Subjective Results.

Results	No. of cases	Percentage
Excellent	14	58.3
Good	5	20.8
Fair	4	16.7
Poor	1	4.2

Post-operative complications: Two of the patients developed superficial skin infections, which were treated with daily dressings and appropriate antibiotics according to the culture and sensitivity reports. All the infections subsided on the above said treatment. We had one patient with ankle stiffness, probably due to the lack of compliance to the advised physiotherapy at home after discharge of the patient. Ankle stiffness ranged from restriction of ankle movement from 20-40%.

DISCUSSION

Fractures of the distal tibia were among the most difficult fractures to treat effectively. The status of the soft tissues, the amount of comminution sustained at the time of injury affect the treatment modality and the long term clinical results. The primary goal of operative treatment is to anatomically align the fractures fragments while providing enough stability to allow early motion. This is achieved with minimal bony and soft tissue devascularisation in order to decrease the complications resulting from the treatment. Gao *et al.* studied 32 adult patients with very short metaphyseal fragments in fractures of distal treated with a polyaxial locking system. The polyaxial locking system shown results of 87.3% excellent to good functional out come with American Orthopedic Foot and Ankle Society score which offer more fixation versatility, may be a reasonable treatment option for distal tibia fracture with very short metaphyseal segments.^[12] Hazarika *et al.* studied a series of 20 patient of distal tibial fracture treated using locking compression plates through MIPPO technique.

This provided 87.5% of good to excellent results. Fractures were classified according to the AO system and performed as scored stage surgery after sterilization with external fixators primarily.^[13] At the end of the our study of 24 patients treated 14(58.3%) patients had excellent outcome, 5(20.8%) had good results, 4(16.7%) had fair outcome and 1(4.2%) had a poor result. Cory collinge *et al.* had an average union of 21 weeks^[14] and Abid mushtaq *et al* had an average of 22 weeks.^[15] Our study had an average fracture union of 16 weeks which were comparable with studies conducted using the locking compression plates.

CONCLUSION

According to this study, 24 patients with fractures of the distal tibia which have undergone closed reduction through MIPPO techniques and application of the locking compression plates states that this technique has resulted in strong and effective stabilization of these fractures. It provides great stability and allows early range of movement at ankle. The closed reduction resulted in rapid union, because it preserves the blood supply to the fragment and helps to achieve near normal anatomical reduction of the fracture.

REFERENCE

1. Michael Sirkin, Roy Sanders. The treatment of pilon fractures. *Clinic Orto*, 2001; 32(1): 91-102.
2. Oni OO, Stafford H, Gergg PJ. A study of diaphyseal fracture repair using tissue isolation techniques. *Injury*, 1992; 23: 467-70.
3. Russell TA. Fractures of the tibia and fibula. In: Rockwood CA, Green DP, Buckolz RW, Heckman JD, editors. *Fractures in adults*. 4th ed. Philadelphia Lippincott; 1996, 2139-57.
4. Surgical Treatment of Nonarticular Distal Tibia Fractures Asheesh Bedi, MD, T. Toan Le, MD and Madhav A. Karunakar, MD 2006 by the American Academy of Orthopaedic Surgeons
5. Shrestha D, Acharya BM, Shrestha PM. Minimal invasive plate osteosynthesis with locking compression plate for distal diaphyseal tibia fracture. *Kathmandu Univ Med J* 2011; 34(2): 62-8.
6. Hasenboehler E, Rikli D, Babst R. Locking compression plate with Minimal invasive plate osteosynthesis in diaphyseal and distal tibial Fracture: a retrospective study of 32 patients. *Injury*, 2007; 38: 365-370.
7. Namazi H, Mozaffarian K. Awful considerations with LCP Instrumentation: a new pitfall. *Arch Orthop Trauma Surg*, 2007; 127: 573-575.
8. Egol KA, Kubiak EN, Fulkerson E, Kummer FJ, Koval KJ. Biomechanics of locked plates and screws. *J Orthop Trauma*, 2004; 18:488-493.
9. Kaab MJ, Frenk A, Schmeling A, Schaser K, Schutz M, Haas NP. Locked internal fixator: sensitivity of screw/plate stability to the correct Insertion angle of the screw. *J Orthop Trauma*, 2004; 18: 483-487.
10. Farouk O, Krettek C, Miclau T *et al.* Minimally invasive plate osteosynthesis and vascularity: preliminary results of a cadaveric injection study. *Injury*, 1999; 30: 591-8.
11. Borelli J, Prickett W, Song E *et al.* Extra osseous blood supply of the tibia and the effects of different plating techniques: human cadaveric study. *J Orthop Trauma*, 2002; 16: 691-5.
12. Gao H, Zhang CQ, Luo CF, Zhou ZB, Zeng BF. Fractures of the distal treated with polyaxial locking plating. *Clin Orthop Relat Res*. Mar., 2009; 467(3): 831-7. Epub 2008 Aug 22.
13. Hazarika S, Chakravarthy J, Cooper J. Minimally invasive locking plate Osteosynthesis for fractures

- of the distal tibia – Injury, 2006; 37(9): 877-87.
Epub 2006 Aug 8.
14. Cory Collinge MD, Mark Kuper DO, Kirk Larson, RNFA, and Robert Protzman MD. Minimally Invasive Plating of High-Energy Metaphyseal Distal Tibial Fractures. J Orthop Trauma, 2007; 21: 355-361.
 15. Abid Mustaq, Rizwan Shahid, Muhammad Asif, Mohammad Maqsood. Distal Tibial fracture fixation with locking compression plate (LCP) using minimally invasive percutaneous osteosynthesis (MIPO) technique. Eur J Trauma Emerg Surg 2008.
 16. Ovadia DN, Beals RK, Fracture of tibial plafond. J Bone Joint Surg 1986; 68(A).